



OAT PROTEIN: HEALTH BENEFITS AND PRODUCT APPLICATIONS

Innovating to Meet Nutrition, Health,
and Wellness Needs Every Day

To learn more about Tate & Lyle ingredients and innovations,
please visit www.foodnutritionknowledge.info
and www.tateandlyle.com.

proatein[®]
Oat Protein

MAKING FOOD EXTRAORDINARY

TATE & LYLE



PROTEIN INTAKE RECOMMENDATIONS AND NEEDS

Protein as a macronutrient

Protein is required to meet nutritional needs and to support health and well-being.¹ This macronutrient is needed to meet human nitrogen requirements and provide indispensable amino acids (also known as essential amino acids). Amino acids are classified as those that cannot be synthesised by the body (indispensable or essential) and those that the body can synthesise (dispensable or nonessential).² However, some are also conditionally indispensable, becoming essential under specific pathological or physiological conditions.² Protein is an important structural and functional component of organs, muscles, biological fluids, and hormones.^{2,3}

Tate & Lyle's PrOatein[®] Oat Protein, like the oat grain it comes from, contains significant amounts of dietary indispensable BCAAs. PrOatein[®] Oat Protein is a natural* protein concentrate ingredient from oats, which can help meet the fast-growing consumer demand for products containing oats and protein-enriched foods.

Global protein intakes

The 2007 World Health Organization (WHO) Technical Report estimates that the average protein requirement for adult maintenance is 105 mg nitrogen per kg body weight per day, or 0.66 g protein per kg body weight per day.⁴ In countries with lower average income, 3% of total dietary energy is from animal protein sources; 11% is from roots and tubers; 6% is from pulses, nuts, and oilseeds; and the remainder is primarily from cereal-based staple foods.⁵ Although the production of livestock

has increased in developing countries, the consumption of protein in these countries remains inadequate.⁵ Additionally, the protein consumed is generally low quality, lacking some of the amino acids required for proper growth and development.⁵

Not every protein source provides all of the amino acids needed for growth and development, and not all protein is equally bioavailable from food sources; however, bioavailability can change with food processing.⁶⁻⁸

Protein quality

Protein digestibility, a component of protein quality, determines the amount of ingested amino acids that are available to the body after digestion and absorption.⁷ Ileal digestibility is the current recommended method for determination of dietary amino acid digestion (further described on the next page), and high levels of digestibility are characteristic for animal proteins and certain purified and concentrated vegetable proteins.^{4,7}

The Food and Agriculture Organization (FAO) of the United Nations Expert Consultation on Protein Quality Evaluation, in conjunction with the WHO, reviewed protein quality assessment and the amino acid scoring.⁹ The 1991 Consultation Report concluded that the Protein Digestibility-Corrected Amino Acid Score (PDCAAS) method, which adjusts amino acid content of the protein source by faecal digestibility correction, was the most suitable evaluation of protein quality for humans.^{2,9-11} This method compares the indispensable amino acid content

- Dietary protein plays essential roles in the body, both structurally (muscle and bone) and functionally (biochemical processes, neurotransmission, circulation).
- The branched-chain amino acids (BCAAs) leucine, isoleucine, and valine stimulate and support muscle protein synthesis (MPS), a process that is important for maintaining muscle structure and function, and are critical for muscle growth and repair at all ages.
- Meta-analyses and clinical studies indicate some favourable effects of higher protein versus lower protein diets on health outcomes including adiposity and cardiovascular risk factors.
- It is recognised that dietary proteins may induce satiety in humans and thus may help to maintain body weight.
- PrOatein[®] Oat Protein contains significant amounts of indispensable BCAAs and can serve as a vegan-friendly protein source for a wide-variety of food applications and nutritional supplements.

*Labeling and/or claims may vary by country. Prospective purchasers are advised to conduct their own tests, studies, and regulatory review to determine the fitness of Tate & Lyle products for their particular purposes, product claims, or specifications.

PrOatein® Oat Protein is a natural protein concentrate ingredient from oats, which can help meet the fast-growing consumer demand for products containing oats and protein-enriched foods.



of the protein source to a reference value for each amino acid based on nutritional needs and corrects for the digestibility of the protein; this method is also recommended by the Codex Committee on Vegetable Proteins.¹¹

An update was published in 2013 and recommended that true ileal digestibility of amino acids from protein sources be accounted for rather than the overall faecal digestibility of a protein.¹ The updated report also includes the adjusted amino acid adequacy reference value recommendations by age group, Table 1.

FAO/WHO now considers the digestible indispensable amino acid score (DIAAS), adjusted for ileal digestibility, to be a replacement for the PDCAAS.^{1, 2, 7, 10} However, PDCAAS is still widely used.

PrOatein® Oat Protein meets nearly all of the FAO 2013 amino acids pattern requirements for adults except for lysine, the limiting amino acid in oat protein. PrOatein® Oat Protein contains at least 10% more sulfur amino acids (SAA) and tryptophan than pea protein; more SAAs and valine than soy protein; and has more of each indispensable amino acid compared to wheat (Figure 1). Combining plant proteins can create a more complete amino acid content in finished food products.

Protein consumption needs for younger populations

Both the quantity and quality of the protein consumed is important to early growth and development as it is well-recognised that protein consumption during this

development window can have effects on long-term health, including influencing body composition.¹² The recommended protein intake for children at two years of age is 0.97 g/kg body weight to ensure adequate intake, which is greater than the minimal amount of protein required for growth.¹² Additionally, the recommended amino acid scoring patterns for young children (aged three years and younger) were revised in the 2013 FAO/WHO report (Table 1).¹ Protein requirements are greater during periods of rapid growth related to the increased demands of growth and increases in body mass and height (see Table 2); furthermore, development of muscles requires additional amino acids.⁴


Table 1
Recommended amino acid scoring patterns^a

Age groups	Scoring pattern (mg/g protein) requirements ^a								
	His	Ile	Leu	Lys	SAA ^b	AAA ^c	Thr	Trp	Val
Child (aged 6 months to 3 years)	20	32	66	57	27	52	31	8.5	43
Older child, adolescent, and adult (>3 years of age)	16	30	61	48	23	41	25	6.6	40

^a Adapted from Chapter 4, Table 5 of the 2013 FAO report

^b Sulfur Amino Acids: Methionine and Cysteine

^c Aromatic Amino Acids: Phenylalanine and Tyrosine

Figure 1Indispensable amino acid contents of selected vegetable protein ingredients^{a,b}^aThree market comparison samples were obtained for soy, pea, and wheat (n=3 for each, n=6 for oat protein).^bThe amino acid and total protein analyses were run at Medallion Laboratories (Minneapolis, MN) via AOAC methods 994.12 and 992.15.

PROTEIN CONSUMPTION AND HEALTH BENEFITS

Increased protein consumption benefits for older populations

Current and emerging evidence suggests that older adults (aged 65 years and older) may need more dietary protein than younger adults to support good health, promote recovery from illness, prevent age-related muscle loss, and maintain muscle functionality.^{13,14} Ageing muscle is less sensitive to the presence of amino acids and may require higher quantities of protein to stimulate muscle protein synthesis (MPS) and accrue muscle proteins.¹⁵ Older adults also may need more protein to offset conditions associated with chronic and acute diseases.^{13,16,17} A recent systematic review examined the health effects of protein intake in healthy elderly adults and found that optimal protein intake is likely higher than the current estimated average requirement (EAR) based on nitrogen balance studies.¹⁶

The PROT-AGE Study Group, an international group of experts representing a wide-range of clinical and research areas, including geriatrics, internal

medicine, endocrinology, nutrition, exercise physiology, gastroenterology, and nephrology, was appointed by the European Union Geriatric Medicine Society, in cooperation with other scientific organisations, to review dietary protein needs with ageing.¹³ The PROT-AGE Study Group published evidence-based recommendations in 2013 for optimal protein intake by older adults recommending an average intake of at least 1.0-1.2 g/kg

body weight per day for people greater than 65 years of age to maintain and regain lean body mass and function.¹³ Additionally, they recommended higher intakes (greater than 1.2 g/kg body weight per day) for those who are physically active. These increased intake targets are recommended due to the reduced ability to use available protein associated with advanced age in order to prevent loss of function in older adults.¹³

Table 2
Minimum protein requirements (g/kg/day)⁴

Age	Maintenance	Growth	Total Need
Birth to 6 months	0.66	0.46	1.12
1 to 2 years	0.66	0.20	0.86
3 to 10 years	0.66	0.07	0.73
11 to 14 years	0.66	0.07	0.73
15 to 18 years	0.66	0.04	0.70
18 years and older	0.66	0.00	0.66

Supplementation with branched-chain amino acids, like those abundant in PrOatein® Oat Protein, has been shown to improve strength and physical function and to increase muscle mass.



Sarcopenia

Sarcopenia, the loss of fat-free mass (FFM) during age-related muscle wasting, can compromise the functional abilities of the elderly.^{15,17} BCAAs have been shown to attenuate muscle wasting,¹⁸ which is important in the prevention of sarcopenia.¹⁹ Dietary leucine may attenuate age-related loss in muscle mass and strength, and may be important in preserving lean muscle mass during ageing.¹⁸

Furthermore, emerging evidence suggests that the elderly specifically may benefit from distributing protein intake evenly throughout the day to promote an optimal per meal stimulation of MPS.¹⁵ Leucine is thought to play a central role in mediating mRNA translation for MPS, and it is recommended that sufficient leucine is provided with dietary protein intake at each meal for the elderly population.¹⁵ Leucine, isoleucine, and valine are BCAAs that are abundant in PrOatein® Oat Protein and are known to influence MPS.

Muscle growth and maintenance

Dietary protein directly contributes to the maintenance of FFM.²⁰⁻²¹ Maintaining skeletal muscle mass is important in conditions such as obesity, hyperlipidemia, cardiovascular disease (CVD), and type 2 diabetes because of the metabolic function of muscle in the body.²²⁻²⁶

The ingestion of protein-containing meals results in stimulation of MPS.¹⁵ In young adults (aged 18 to 30 years old), supplementation with a leucine-rich, high-quality protein can augment exercise-induced muscle mass development and lead to strength gains to a greater extent compared to other protein sources.^{15, 21, 27-28} Additionally, supplementation with leucine-rich amino acid mixtures has been shown to improve strength and physical function and to increase muscle mass in the elderly.^{15, 29}

When combined with exercise, protein ingestion helps to increase muscle synthesis, thus assisting in muscle protein accretion.¹⁵

Physical activity can increase the demands for protein consumption.⁴ The indispensable BCAA leucine is a direct regulator of MPS through activation of the mTOR pathway, a cell signal known to stimulate muscle protein production.^{15, 18}

Furthermore, the BCAAs leucine, isoleucine, and valine compose 14-18% of the total amino acids found in skeletal muscle protein and all of these amino acids are required for maintaining muscle health.¹⁸

Leucine, along with isoleucine and valine, has been shown to stimulate MPS when administered in a protein-containing beverage in young men.³⁰ After exercise, the repeated ingestion of 20 g of high-quality protein containing leucine also has been shown to provide maximal muscle building stimulus during the recovery period.³¹

Satiety and weight management

Obesity continues to be a major public health concern globally.³² While energy balance is key to weight management, scientific research suggests that a diet rich in high-quality protein is one dietary strategy to aid in acute postprandial satiety and thus help with weight maintenance.^{14, 20, 25-26, 33-35} High-protein diets have been successful at preserving lean body mass during weight loss,^{14, 34, 36} and diets using meal replacements that provide higher protein with moderate fat have been shown to assist in weight maintenance.³⁷ In the context of calorie reduction, a recent review found that a protein intake of 0.8-1.2 g/kg body weight per day is sufficient to sustain satiety, energy expenditure, and FFM independent of dietary carbohydrate content.²⁰

The protein content of the diet has long been recognised for its effect on food intake because high protein diets may promote satiety through reduced calorie intake, increase the postprandial perception of satiety, and stimulate the endocrine hormones in the gastrointestinal tract known to increase satiety.^{14, 25, 33, 35, 38-42} Some research supports that a higher protein intake (25% of total energy) is effective for appetite control and satiety in overweight and obese men during hypocaloric-induced weight loss.⁴³ Additionally, when overweight and obese teen girls consumed a high-protein breakfast meal (35 g protein), there was a

While energy balance is key to weight management, scientific research suggests that a diet rich in high quality protein is one dietary strategy to aid in acute postprandial satiety and thus help with weight maintenance.



significant reduction in four-hour cravings for savory foods.⁴⁰ Also, it has been shown that consumption of an afternoon yoghurt snack (containing 14 g protein) versus crackers (containing 0 g protein) versus chocolate (containing 2 g protein) can increase the time to initiation of the next meal and decrease caloric intake in healthy women.⁴¹ Gut hormones work collectively in response to meal consumption to help regulate food intake.^{44, 45}

Changes in perception of postprandial hunger, satiety, and food intake are positively associated with meals containing proteins.^{43, 46} The ratio of macronutrients in a study of isocaloric meals demonstrated that a low-

carbohydrate/high-protein diet (65.0% calories from protein, 17.2% calories from carbohydrate) significantly increased peptide YY (PYY) and decreased hunger in humans when compared to a high-carbohydrate/low-protein diet (17.8% calories from protein and 64.5% calories from carbohydrates). This effect was seen in both lean and overweight individuals.³³ Although more research is needed in this area, increasing protein intake may be one way to help manage food intake and may help to maintain a healthy body weight.


Cardiovascular benefits

Research suggests the consumption of plant proteins may have protective effects against chronic diseases and may contribute to

decreased circulating cholesterol levels.⁴⁷⁻⁵² Plant protein food sources are generally low in fat and can displace other dietary sources of protein that provide higher amounts of fat.^{20, 52} Furthermore, it is well known that diets high in fruits, whole grains, and vegetables are associated with a lower rate of CVD.^{20, 48, 53, 54}

Prospective cohort studies support that high consumption of plant-based foods is also associated with a significantly lower risk of coronary artery disease and stroke.^{20, 48} Higher protein diets in general may improve blood pressure,^{48, 54, 55} triglyceride levels, and reduce adiposity according to a recent systematic review of 74 randomised clinical trials.³⁴

Figure 2
Indispensable amino acid contents of PrOatein® Oat Protein^a



^a Average of six independently tested lots of PrOatein® Protein.

* PrOatein® Oat Protein has a minimum 50% protein content specification, with a typical range of 52-56% protein content (on a dry matter basis).

** Labeling and/or claims may vary by country. Prospective purchasers are advised to conduct their own tests, studies, and regulatory review to determine the fitness of Tate & Lyle products for their particular purposes, product claims, or specifications.

PROATEIN® OAT PROTEIN CONCENTRATE

In addition to its typical 55%* protein content, PrOatein® Oat Protein contains:

- Typically 16-19% naturally occurring oat oil composed of:
 - Primarily palmitic, oleic, and linoleic fatty acids
 - A high concentration of monounsaturated fatty acids (oleic acid) and a high amount of polyunsaturated fatty acids (mainly omega-6)
 - Micronutrients with antioxidant properties, including vitamin E (tocopherols) and avenanthramides
 - No cholesterol, since this is a plant-based protein
- Naturally-occurring oat maltodextrins (approx. 20-24%)
- A small amount (2%) of oat beta glucan soluble fibre
- PrOatein® Oat Protein has a great taste compared to other isolated proteins
- PrOatein® Oat Protein is a natural protein concentrate ingredient made from oats that can help meet the fast-growing consumer demand for protein-enriched foods and products containing oats
- PrOatein® Oat Protein is vegan-friendly

PROATEIN® OAT PROTEIN CONCENTRATE

Amino acid content

PrOatein® Oat Protein** is made from Swedish oats and is vegetarian- and vegan-friendly. PrOatein® Oat Protein contains many of the amino acids needed for growth and development and has an abundance of leucine, isoleucine, and valine, the BCAAs known to be involved in muscle protein synthesis (Figure 2). As for oats in general, lysine is the limiting amino acid in oat protein. Combining oat protein with other plant proteins, such as pea protein, can create a more complete amino acid content in finished food products while still remaining vegetarian and vegan. When adjusting for a 91% protein digestibility (based on rolled oats⁹), the PDCAAS for PrOatein® Oat Protein is 46%.

PrOatein® Oat Protein has a great taste compared to other isolated proteins such as soy protein. PrOatein® Oat Protein is a natural protein concentrate from oats that can help meet the fast-growing consumer demand for protein-enriched foods.

Use PrOatein® Oat Protein in a wide-variety of food applications and nutritional supplements

Oat-based products and protein-fortified foods are becoming increasingly popular in the marketplace. By using PrOatein® Oat Protein, a wide-variety of food applications and nutritional supplements can provide the benefits of added protein that consumers desire while delivering great taste that consumers expect.

PrOatein® Oat Protein performs well in many food applications including bars, breakfast cereals and porridges, breads, cookies, supplements, and sports nutrition beverages. With its mild cereal taste, PrOatein® Oat Protein also makes a perfect base for bar fillings. PrOatein® Oat Protein concentrate is a natural clean-label ingredient made from oats without the addition of chemicals or solvents.





To learn more about Tate & Lyle ingredients and innovations as well as health benefits and relevant research, please visit www.foodnutritionknowledge.info and www.tateandlyle.com.

INNOVATING TO MEET NUTRITION, HEALTH, AND WELLNESS NEEDS EVERY DAY

Conclusions

Protein is an important part of the diet and plays an essential role in the structural and functional components of the body. PrOatein® Oat Protein, like the oat grain it comes from, contains significant amounts of dietary indispensable amino acids, particularly the BCAAs leucine, isoleucine, and valine, which are important for muscle growth and repair in people of all ages. Oat protein contains more SAA and tryptophan than pea protein; more SAA and valine than soy protein; and has more indispensable amino acids than wheat protein. PrOatein® Oat Protein also has a great taste compared to other isolated plant proteins.

Ongoing research indicates that dietary proteins may induce satiety and could thus help with maintenance of body weight. Consequently, supplementing commonly-consumed foods with protein may help increase satiety and possibly decrease total energy intake.

A commitment to innovation

Tate & Lyle, a global leader in wellness innovation, is committed to delivering innovative ingredients that can be incorporated into great-tasting foods to help consumers meet their nutrition, health, and wellness needs every day. That is because Tate & Lyle invests heavily in innovation and research and in developing ingredients that can be incorporated into a wide-variety of food and beverage solutions. Teams of food and nutrition scientists are continuously innovating, researching, and testing ingredients that will meet current and future health and nutrition needs.

At the same time, Tate & Lyle has a robust market research program designed to provide the necessary insights on consumer preferences around the world. The research program allows Tate & Lyle to customise its offerings and provide tailor-made solutions in local and regional markets.

Better-for-you ingredients for health and wellness

In response to global public health efforts calling for people to reduce calories and sodium and increase fibre intakes, Tate & Lyle offers a number of innovative ingredient solutions that meet these needs.

To learn more about Tate & Lyle ingredients and innovation as well as health benefits and relevant research, please visit www.foodnutritionknowledge.info and www.tateandlyle.com.

REFERENCES

- 1 Consultation FE. Dietary protein quality evaluation in human nutrition. FAO Food and Nutrition Paper. 2013:1-66.
- 2 Boye J, Wijesinha-Bettoni R, Burlingame B. Protein quality evaluation twenty years after the introduction of the protein digestibility corrected amino acid score method. *Br J Nutr.* 2012;108 Suppl 2:S183-211.
- 3 Fleddermann M, Fechner A, Rossler A, Bahr M, Pastor A, Liebert F, Jahreis G. Nutritional evaluation of rapeseed protein compared to soy protein for quality, plasma amino acids, and nitrogen balance—a randomized cross-over intervention study in humans. *Clin Nutr.* 2013;32:519-526.
- 4 World Health Organization. Proteins and Amino Acids in Human Nutrition. WHO/FAO/UNU Expert Consultation. World Health Organ Tech Rep. 2007;935.
- 5 Schonfeldt HC, Gibson Hall N. Dietary protein quality and malnutrition in Africa. *Br J Nutr.* 2012;108 Suppl 2:S69-76.
- 6 Potier M, Tome D. Comparison of digestibility and quality of intact proteins with their respective hydrolysates. *J AOAC Int.* 2008;91:1002-1005.
- 7 Tome D. Digestibility issues of vegetable versus animal proteins: protein and amino acid requirements—functional aspects. *Food Nutr Bull.* 2013;34:272-274.
- 8 Tome D. Criteria and markers for protein quality assessment—a review. *Br J Nutr.* 2012;108 Suppl 2:S222-229.
- 9 Protein quality evaluation. Joint FAO/WHO. FAO Food Nutr Pap 1991;51:1066.
- 10 Gilani GS. Background on international activities on protein quality assessment of foods. *Br J Nutr.* 2012;108 Suppl 2:S168-182.
- 11 Lewis JL. The regulation of protein content and quality in national and international food standards. *Br J Nutr.* 2012;108 Suppl 2:S212-221.
- 12 Michaelsen KF, Greer FR. Protein needs early in life and long-term health. *Am J Clin Nutr.* 2014;99:718S-722S.
- 13 Bauer J, Biolo G, Cederholm T, Cesari M, Cruz-Jentoft AJ, Morley JE, et al. Evidence-based recommendations for optimal dietary protein intake in older people: a position paper from the PROT-AGE Study Group. *J Am Med Dir Assoc.* 2013;14:542-559.
- 14 Paddon-Jones D, Leidy H. Dietary protein and muscle in older persons. *Curr Opin Clin Nutr Metab Care.* 2014;17:5-11.
- 15 Breen L, Phillips SM. Skeletal muscle protein metabolism in the elderly: Interventions to counteract the ‘anabolic resistance’ of ageing. *Nutr Metab (Lond).* 2011;8:68.
- 16 Pedersen AN, Cederholm T. Health effects of protein intake in healthy elderly populations: a systematic literature review. *Food Nutr Res.* 2014;58.
- 17 Beasley JM, Shikany JM, Thomson CA. The role of dietary protein intake in the prevention of sarcopenia of aging. *Nutr Clin Pract.* 2013;28:684-690.
- 18 Manders RJ, Little JP, Forbes SC, Candow DG. Insulinotropic and muscle protein synthetic effects of branched-chain amino acids: potential therapy for type 2 diabetes and sarcopenia. *Nutrients.* 2012;4:1664-1678.
- 19 Ribeiro SM, Kehayias JJ. Sarcopenia and the analysis of body composition. *Adv Nutr.* 2014;5:260-267.
- 20 Martens EA, Westerterp-Plantenga MS. Protein diets, body weight loss and weight maintenance. *Curr Opin Clin Nutr Metab Care.* 2014;17:75-79.
- 21 Phillips SM. Dietary protein requirements and adaptive advantages in athletes. *Br J Nutr.* 2012;108 Suppl 2:S158-167.
- 22 Dideriksen K, Reitelseder S, Holm L. Influence of amino acids, dietary protein, and physical activity on muscle mass development in humans. *Nutrients.* 2013;5:852-876.
- 23 Wolfe RR. The underappreciated role of muscle in health and disease. *Am J Clin Nutr.* 2006;84:475-482.
- 24 Hu FB. Plant-based foods and prevention of cardiovascular disease: an overview. *Am J Clin Nutr.* 2003;78:544S-551S.
- 25 Leidy HJ. Increased dietary protein as a dietary strategy to prevent and/or treat obesity. *Mo Med.* 2014;111:54-58
- 26 Keller U. Dietary proteins in obesity and in diabetes. *Int J Vitam Nutr Res.* 2011;81:125-133.
- 27 Hartman JW, Tang JE, Wilkinson SB, Tarnopolsky MA, Lawrence RL, Fullerton AV, Phillips SM, et al. Consumption of fat-free fluid milk after resistance exercise promotes greater lean mass accretion than does consumption of soy or carbohydrate in young, novice, male weightlifters. *Am J Clin Nutr.* 2007;86:373-381.
- 28 Millward DJ. Knowledge gained from studies of leucine consumption in animals and humans. *J Nutr.* 2012;142:2212S-2219S.
- 29 Scognamiglio R, Testa A, Aquilani R, Dioguardi FS, Pasini E. Impairment in walking capacity and myocardial function in the elderly: is there a role for nonpharmacologic therapy with nutritional amino acid supplements? *Am J Cardiol.* 2008;101:78E-81E.
- 30 Churchward-Venne TA, Breen L, Di Donato DM, Hector AJ, Mitchell CJ, Moore DR, et al. Leucine supplementation of a low-protein mixed macronutrient beverage enhances myofibrillar protein synthesis in young men: a double-blind, randomized trial. *Am J Clin Nutr.* 2014;99:276-286.
- 31 Areta JL, Burke LM, Ross ML, Camera DM, West DW, Broad EM, et al. Timing and distribution of protein ingestion during prolonged recovery from resistance exercise alters myofibrillar protein synthesis. *J Physiol.* 2013;591:2319-2331.
- 32 World Health Organization. Obesity and Overweight Fact Sheet; Version No. 311. August 2014.
- 33 Batterham RL, Heffron H, Kapoor S, Chivers JE, Chandarana K, Herzog H, et al. Critical role for peptide YY in protein-mediated satiation and body-weight regulation. *Cell Metab.* 2006;4:223-233.
- 34 Santesso N, Akl EA, Bianchi M, Mente A, Mustafa R, Heels-Ansdell D, Schunemann HJ. Effects of higher- versus lower-protein diets on health outcomes: a systematic review and meta-analysis. *Eur J Clin Nutr.* 2012;66:780-788.
- 35 Veldhorst M, Smeets A, Soenen S, Hochstenbach-Waelen A, Hursel R, Diepvens K, Lejeune M, Luscombe-Marsh N, Westerterp-Plantenga M. Protein-induced satiety: effects and mechanisms of different proteins. *Physiol Behav.* 2008;94:300-307.
- 36 Soenen S, Martens EA, Hochstenbach-Waelen A, Lemmens SG, Westerterp-Plantenga MS. Normal protein intake is required for body weight loss and weight maintenance, and elevated protein intake for additional preservation of resting energy expenditure and fat free mass. *J Nutr.* 2013;143:591-596.
- 37 Soeliman FA, Azadbakht L. Weight loss maintenance: A review on dietary related strategies. *J Res Med Sci.* 2014;19:268-275.
- 38 Davidenko O, Darcel N, Fromentin G, Tome D. Control of protein and energy intake—brain mechanisms. *Eur J Clin Nutr.* 2013;67:455-461.
- 39 Journal M, Chaumontet C, Darcel N, Fromentin G, Tome D. Brain responses to high-protein diets. *Adv Nutr.* 2012;3:322-329.
- 40 Hoertel HA, Will MJ, Leidy HJ. A randomized crossover, pilot study examining the effects of a normal protein vs. high protein breakfast on food cravings and reward signals in overweight/obese “breakfast skipping,” late-adolescent girls. *Nutr J.* 2014;13:80.
- 41 Ortinau LC, Hoertel HA, Douglas SM, Leidy HJ. Effects of high-protein vs. high- fat snacks on appetite control, satiety, and eating initiation in healthy women. *Nutr J.* 2014;13:97.
- 42 Simpson K, Parker J, Plumer J, Bloom S. CCK, PYY and PP: the control of energy balance. *Handb Exp Pharmacol.* 2012;209-230.

- 43 Leidy HJ, Tang M, Armstrong CL, Martin CB, Campbell WW. The effects of consuming frequent, higher protein meals on appetite and satiety during weight loss in overweight/obese men. *Obesity (Silver Spring)*. 2011;19:818-824.
- 44 Stengel A, Tache Y. Interaction between gastric and upper small intestinal hormones in the regulation of hunger and satiety: ghrelin and cholecystokinin take the central stage. *Curr Protein Pept Sci*. 2011;12:293-304.
- 45 Begg DP, Woods SC. The endocrinology of food intake. *Nat Rev Endocrinol*. 2013;9:584-597.
- 46 Smith CE, Mollard RC, Luhovyy BL, Anderson GH. The effect of yellow pea protein and fibre on short-term food intake, subjective appetite and glycaemic response in healthy young men. *Br J Nutr*. 2012;108 Suppl 1:S74-80.
- 47 Clifton PM. Protein and coronary heart disease: the role of different protein sources. *Curr Atheroscler Rep*. 2011;13:493-498.
- 48 Shay CM, Stamler J, Dyer AR, Brown IJ, Chan Q, Elliott P, et al. Nutrient and food intakes of middle-aged adults at low risk of cardiovascular disease: the international study of macro-/micronutrients and blood pressure (INTERMAP). *Eur J Nutr*. 2012;51:917-926.
- 49 Krajcovicova-Kudlackova M, Babinska K, Valachovicova M. Health benefits and risks of plant proteins. *Bratisl Lek Listy*. 2005;106:231-234.
- 50 Jenkins DJ, Wong JM, Kendall CW, Esfahani A, Ng VW, Leong TC, et al. Effect of a 6-month vegan low-carbohydrate ('Eco-Atkins') diet on cardiovascular risk factors and body weight in hyperlipidaemic adults: a randomised controlled trial. *BMJ Open*. 2014;4:e003505.
- 51 Reynolds K, Chin A, Lees KA, Nguyen A, Bujnowski D, He J. A meta-analysis of the effect of soy protein supplementation on serum lipids. *Am J Cardiol*. 2006;98:633-640.
- 52 Anderson JW, Bush HM. Soy protein effects on serum lipoproteins: a quality assessment and meta-analysis of randomized, controlled studies. *J Am Coll Nutr*. 2011;30:79-91.
- 53 Ha V, Sievenpiper JL, de Souza RJ, Jayalath VH, Mirrahimi A, Agarwal A, et al. Effect of dietary pulse intake on established therapeutic lipid targets for cardiovascular risk reduction: a systematic review and meta-analysis of randomized controlled trials. *CMAJ*. 2014;186:E252-262.
- 54 Pedersen AN, Kondrup J, Borsheim E. Health effects of protein intake in healthy adults: a systematic literature review. *Food Nutr Res*. 2013;57.
- 55 Buendia JR, Bradlee ML, Singer MR, Moore LL. Diets Higher in Protein Predict Lower High Blood Pressure Risk in Framingham Offspring Study Adults. *Am J Hypertens*. 2014.

This leaflet is provided for general circulation to the nutrition science and health professional community and professional participants in the food industry, including prospective customers for Tate & Lyle food ingredients. It is not designed for consumer use. The applicability of label claims, health claims and the regulatory and intellectual property status of our ingredients varies by jurisdiction. You should obtain your own advice regarding all legal and regulatory aspects of our ingredients and their usage in your own products to determine suitability for their particular purposes, claims, freedom to operate, labeling or specific applications in any particular jurisdiction. This product information is published for your consideration and independent verification. Tate & Lyle accepts no liability for its accuracy or completeness. Tate & Lyle • 5450 Prairie Stone Parkway, Hoffman Estates, IL 60192 • 1.800.526.5728

SOG1017019